

Evaluation of 21 Brassica microgreens growth and nutritional profile grown under different red, blue and green LEDs combination

Khaled Y. Kamal^{1,2*}, Ahmed A. El-Tantawy³, Dina Abdel Moneim⁴, Asmaa Abdel Salam¹, Naglaa Qabil¹, Salwa, M. A. I. Ash-shormillesy¹, Ahmed Attia¹, Mohamed A. S. Ali⁵, Raúl Herranz⁶, Mohamed A. El-Esawi^{7,8}, Amr A. Nassrallah⁹

¹Agronomy Department, Faculty of Agriculture, Zagazig University, Zagazig, Sharqia, 44511, Egypt

²Department of Molecular Biology, Centre of the Region Haná for Biotechnological and Agricultural Research, Faculty of Science, Palacký University, Olomouc, Czech Republic.

³Ornamental Horticulture Department, Faculty of Agriculture, Cairo University, Cairo, 12613, Egypt

⁴Department of Plant production (Genetic branch), Faculty of Environmental and Agricultural Sciences, Arish University, Egypt

⁵Plant Pathology Department, Faculty of Agriculture, Zagazig University, Sharqia, 44511, Egypt

⁶Centro de Investigaciones Biológicas (CSIC), Ramiro de Maeztu 9, 28040 Madrid, Spain

⁷Botany Department, Faculty of Science, Tanta University, 31527 Tanta, Egypt

⁸Sainsbury Laboratory, University of Cambridge, Cambridge, United Kingdom

⁹Biochemistry Department, Faculty of Agriculture, Cairo University, Cairo, Egypt

***Corresponding author:**

Dr. Khaled Youssef Kamal

ORCID: <https://orcid.org/0000-0002-6909-8056>

Faculty of Agriculture, Zagazig University

Al Wehda Al Zraaia St, AZ Zagazig, 44519, Egypt

Tel: +20 55 2245274 Fax: +20 55 2221688

Email: kymostafa@agri.zu.edu.eg

35

36 Abstract

37 Microgreens are rich functional crops with valuable nutritional elements that have health benefits
 38 when used as food supplants. Growth characterization, nutritional composition profile of 21
 39 varieties representing 5 species of the *Brassica* genus as microgreens were assessed under light-
 40 emitting diodes (LEDs) conditions. Microgreens were grown under four different LEDs ratios (%)
 41 (R₈₀:B₂₀, R₂₀:B₈₀, R₇₀:G₁₀:B₂₀, and R₂₀:G₁₀:B₇₀). Results indicated that supplemental lighting with
 42 green LEDs (R₇₀:G₁₀:B₂₀) enhanced vegetative growth and morphology, while blue LEDs (R₂₀:B₈₀)
 43 increased the mineral composition and vitamins content. Interestingly, combining the nutritional
 44 content with the growth yield to define the optimal LEDs setup, we found that the best lighting to
 45 promote the microgreen growth was supplying the green LEDs combination (R₇₀:G₁₀:B₂₀).
 46 Remarkably, under this proper conditions, Kohlrabi purple, Cabbage red, Broccoli, Kale Tucsan,
 47 Komatsuna red, Tatsoi, and Cabbage green had the highest growth and nutritional content profile as
 48 microgreens which being a health-promoting in a diet support strategy required for the human
 49 health under certain isolated of limited food conditions.

50

51 **Keywords:** Brassicaceae; Functional Crops; Light Emitting Diodes; Microgreens; Nutritional
 52 quality

53

54

55

56

57

58 1. Introduction

59 As the world's population is rapidly growing, with an increasing demand for sustainable sources of
 60 food products such as the rich-nutrient functional crops. Ongoing efforts are aimed to find new
 61 strategies for food production to meet the demands of the growing world population. Recently, the
 62 consumption of microgreens has increased, as a rich-nutrient crop with a high level of nutrition
 63 components concentration contains; vitamins, minerals, and antioxidants compared to mature
 64 greens, which are helpful in filling the nutritional gap challenges [1]. Furthermore, microgreens
 65 being valuable functional crops for their rich-phytonutrients content [2, 3]. Microgreens are a
 66 category of edible salad crops that appearing in many upscale markets and restaurants. They are
 67 harvested at the base of the hypocotyl when the first true leaves start to emerge, generally, the
 68 growth rate is ≤ 21 days after sowing [4, 5]. Despite their small size, they can provide a high
 69 concentration of health-promoting phytochemicals [5]. Commercially greenhouse growers became
 70 more interested in the microgreen for their high market levels [4]. Specifically, microgreens of the
 71 family *Brassicaceae* have become a popular choice due to its easy way for germination and short
 72 growth length and providing wide flavors and colors [5]. *Brassicaceae* microgreens species could be
 73 used as a new ingredient which provides a wide variety of our food [5-7] and valued for containing
 74 significant amounts of cancer-fighting glucosinolates [8]. They are also rich in carotenoids,
 75 especially lutein, zeaxanthin, and β -carotene [9-11]. Thus, brassica microgreens are considered as a
 76 functional food, which serves as a health-promoting or disease preventing supplementals [5, 12]

77 Several strategies were used and developed for providing optimal greenhouse conditions to increase
 78 the microgreen yield. Light emitting diodes (LEDs) is a new light source technology used for
 79 greenhouses facilities and space- limited plant growth chambers [13, 14]. It becomes more
 80 economically viable with high efficiency and low cost, as well as the ability to select light qualities

and intensities [15]. It is reported that crop plants use light for photosynthesis and being responded to the different light intensity, wavelength [16, 17]. Microgreens have a lower demand for photon flux compared to long-cycle crops, thus are ideally adapted to chamber environments. Recently, many studies demonstrated the influence of LEDs (blue or/and red) lighting on the plant vegetative parameters [14, 18, 19] and demonstrated the effect of light quality on the growth of the cultivated plants [8, 20-22]. Nevertheless, a lack of information regarding the combined effect of red and blue and other LEDs lighting such as green light on the plant growth, morphology, and nutrition content profile of microgreens [22, 23]. Furthermore, green light supplies enhance the carotenoid content in mustard microgreens [24].

Although microgreens have been considered as valuable and nutritionally beneficial functional crops, a little is known on the integrity of individual and combined influence of green, red, and blue LEDs on Brassica species microgreens growth and nutritional composition. Therefore, the main purpose of this current study is to define the influence of alternative LEDs light regimens on Brassica species microgreens growth, and nutritional composition and to define which species could serve well as a life support component in many cases. We explore the impact of different four LEDs lighting ratio (Red, Blue, and Green) on 21 Brassica microgreens growth and nutritional profile.

2. Material and Methods

2.1.Plant Materials and Growth chamber environment

Twenty-one varieties of microgreens representing 5 species of Brassica genus of the *Brassicaceae* family (Table 1) were grown in greenhouse chambers in a collaborated study between the Faculty of Agriculture in both Zagazig University and Cairo University. We used the recommended soil and fertilization properties as reported by [5]. About 10-25 g of seeds, varying based on the seed index of each variety, (Table 1) were sown in peat moss in Rockwool tray in a controlled conditions

greenhouse (3 trays per each variety for 3 replicates), cultivated under relative humidity (RH), and carbon dioxide (CO₂) concentration of 70%, and 500 $\mu\text{mol.mol}^{-1}$, respectively. Each day, 100 ml of CaCl₂ solution was added to each tray to further stimulate seedling growth. Once cotyledons were fully reflexed 5 d after sowing, 300 ml of 25% nutrient solution was added to each tray daily until harvest. This experiment was carried out simultaneously in the summer season of 2018 from May to September with as a growth length for each species ranging from 6-12 days (Table 1).

2.2.LEDs lighting treatments

Brassica microgreen plants were grown under LEDs lighting (Light-emitting diode arrays) were provided by four different light quality ratios (%) treatments of red:blue 80:20 and 20:80 (R₈₀:B₂₀, and R₂₀:B₈₀), or red:green:blue 70:10:20, and 20:10:70 (R₇₀:G₁₀:B₂₀, and R₂₀:G₁₀:B₇₀) (Philips GreenPower LED production modules; Koninklijke Philips Electronics, N. V., Amsterdam, The Netherlands), using 0.5 W per LED chip. Each LEDs treatment was carried out in a different room. In the controlled environment greenhouse, the LEDs were placed horizontally, above the bench top, at a height of 50 cm. we adjusted the photosynthetic photon flux density (PPFD) average to 150 $\mu\text{mol.m}^{-2}.\text{s}^{-1}$ that was provided by the fluorescent lamps and bar-type LEDs. This experiment was performed three times replications with the same conditions.

2.3.Harvest, Growth measurements

Microgreen samples were harvested after the growth length for each species (Table1) without seed coats or roots as recommended by [5]. Each replicate used for the measurements consisted of at least 10 grown seedlings. Ten seedlings of each microgreen variety were randomly selected and measured to determine Hypocotyl Length (HL), Leaf Area (LA), for each LEDs treatment. Hypocotyl measurements HL of the harvested seedlings were measured from the tip where the cotyledons split, to the end of the base of the hypocotyl. LA of cotyledons and fully expanded

leaves were measured by LA meter (LI-3100; LI-COR Inc. Lincoln, NE) by recording the average of five scans.

Furthermore, another ten randomly selected seedlings for each variety used to assess both, Fresh weight (FW), and Dry weight percentage (DW%). After FW data were measured, samples were oven dried at 80°C for 72 hours. Then DW data were measured. FW and DW values were used to calculate DW% ($DW\% = (DW/FW \times 100)$).

2.4.Elemental Analysis

Fresh microgreens (50 g FW per each sample) were collected and rinsed 3X using H₂O to remove any surface residue. Dried microgreens (2 g per replicate) were grounded into a fine powder to analysis the elemental composition. , Each of the 21 samples was subjected to acid digestion procedures and quantitative measurements of the following elements: P, K, Ca, Na, Fe, Mn, Cu, and Zn were done using inductively coupled plasma optical emission spectrometry (ICP-OES) following the methods of Huang and Schulte [25]. To assure the accuracy of the method, standard reference materials (Apple leaves, NIST® SRM® 1515, NIST1515, SIGMA, USA, and Spinach leaves, NIST® SRM® 1570a, NIST1570A) were used and evaluated using the same digested procedure. For each ICP-OES analyte, the limit of detection (LOD) and limit of quantification (LOQ), which are a function of the sample mass were determined (Supplementary Table 1)

2.5.Vitamin and Carotenoid concentration analysis

2.5.1. Phylloquinone

Phylloquinone was determined according to a previously reported method by [26]. Under dim light, 0.2 g of dried microgreens were homogenized in 10 mL of H₂O and 0.4 mL of 200 µg/mL menaquinone used as an internal standard. The sample was supplied with 15 mL of 2-

propanol/hexane (3:2 v/v) and were vortexed for 1 min. Then the sample was centrifuged at 1500g at 21°C for 5 min. Then we transferred the upper layer (hexane) into a new glass tube and to dry using a stream of N₂. The residues of the sample were dissolved using 4 mL of hexane. Then, to purify the extract, 1 mL of the dissolved extract was loaded onto preconditioned silica gel columns (4 mL of 3.5% ethyl ether in hexane, followed by 4 mL of 100% hexane). We used 2 mL of hexane to wash the columns. Phylloquinone was eluted with 8 mL of 3.5% ethyl ether in hexane and then evaporated at 40 °C under N₂ flow. Further, it is reconstituted in 2 mL of mobile phase (99% methanol and 1% 0.05 M sodium acetate buffer, pH = 3.0) and is filtered through a 0.22 µm nylon syringe filter (Millipore, Bedford, MA). To detect the phylloquinone, we used a photodiode array detector (DAD) (G1315C, Agilent, Santa Clara, CA) on Agilent 1200 series HPLC system and absorbance wavelength was 270 nm. 20 µL of the extract was injected into the HPLC and being run through a C18 column (201TP, 5 µm, 150 × 4.6 mm, Grace, Deerfield, IL) flowing at the rate of 1 mL/ min. The phylloquinone content was measured according to the internal standard method based on peak areas.

2.5.2. Carotenoids and Tocopherols

To extract both carotenoids and tocopherols, we followed the procedure described by [27] and modified by Xiao et al. [5]. In 15 mL screw-cap glass vial, 0.05 g of dried fine powder was homogenized in 7.5 mL of 1% butylated hydroxytoluene in ethanol and 500 µL of 86.82 µM trans-βapo-8 carotenal as an internal standard was added. 180 µL of 80% KOH was supplied to the mixture and, the vials were capped and placed in a dry bat at 70 °C for 15 min. The vials were removed and being cooled on ice 4°C for 5 min. The mixture was transferred into 15 mL centrifuge tubes supplied with 3.0 mL of deionized water and 3.0 mL of hexane/toluene solution (10:8 v/v). The mixture was carefully vortexed for 1 min and then were centrifuged at 1000g for 5 min. After centrifugation, the upper organic layer was collected into an 8 mL glass culture tube and

immediately placed into a nitrogen evaporator set at 30 °C. on the other hand, the lower layer was extracted with 3.0 mL of hexane/toluene (10:8 v/v). this extraction process was repeated at least four times until the upper layer is colorless. After evaporation, the residue was diluted in 500 µL of mobile phase acetonitrile/ethanol (1:1 v/v), filtered into an HPLC amber vial using nylon filter (0.22 µm, Millipore, Bedford, MA). Subsequently, 20 µL was inoculated for HPLC analysis. Carotenoid and tocopherol concentrations were measured using isocratic reverse-phase high-performance liquid chromatography (RP-HPLC). Absorbance was measured at 290 nm (for tocopherols) and 450 nm (for carotenoids).

2.5.3. Ascorbic Acid

Total ascorbic acid (TAA) was assessed spectrophotometrically according to [28]. 3g fresh microgreens were homogenization in 10 mL of ice-cold 5% metaphosphoric acid (w/v) at 4°C at 15000 rpm for 1 min. The homogenized then centrifuged at 7000g for 20 min at 4°C. The supernatant was filtered through Whatman 4# filter paper. TAA was measured spectrophotometrically at 525 nm. Concentrations of TAA was calculated from an L-ascorbic acid standard curve.

2.6. Clustering hierarchical analysis

In order to extrapolate the similarities and the dissimilarities among the 21 microgreens in growth and nutritional assessment, hierarchical cluster analysis was performed using the normalized data set using *class Orange clustering hierarchical* using *ORANGE version 3.7* [29].

2.7. Statistical analysis

The experiment was laid out in a randomized block design in a factorial arrangement with LEDs (four levels) and Microgreens (Twenty-one varieties) for three different biological replicates. Data

were collected and analyzed according to [30]. SPSS_{v.22} software was used to analyze the variance of differences using ANOVA test statistically followed by LSD analysis. The degree of freedom was followed as $P \leq 0.05$, $P \leq 0.01$, and $P \leq 0.001$ considers the statistical significance and represents as *, **, *** respectively.

3. Results.

3.1. The influence of LEDs on microgreens growth, nutritional profile

In the present work, four different LEDs lighting ratio (%) treatments of R₈₀:B₂₀, R₂₀:B₈₀, R₇₀:G₁₀:B₂₀, and R₂₀:G₁₀:B₇₀ were implemented. Growth parameters of the 21 varieties were analyzed (Fig. 1). Results revealed that those microgreens are grown under the R₇₀:G₁₀:B₂₀ had the highest growth and morphology targeted parameter, while the lowest growth parameters were observed under R₂₀:B₈₀ (Figure 1). The Hypocotyl length (HL) and leaf area (LA) of the microgreens were significantly elongated in plants grown under R₇₀:G₁₀:B₂₀ compared to those grown under R₈₀:B₂₀, R₂₀:B₈₀, and R₂₀:G₁₀:B₇₀, respectively (Figure1)., Fresh weight (FW) and Dry weight % (DW%) of those microgreens grown under R₇₀:G₁₀:B₂₀ treatment showed the highest values; on average; 0.4g (FW), and 6.27 %(DW%). Indicating that R₇₀:G₁₀:B₂₀ combination induces an increase in all studied growth and morphology parameters in comparison with the other LEDs lighting treatments (Figure1).

Considering that R₇₀:G₁₀:B₂₀ LEDs lighting combination has an impact in targeted growth parameters, we investigated whether it has a functional influence on the nutritional composition profile by conducting an ICP analysis of macro and microelements from 21 varieties Brassicaceae microgreen using lowest growth enhancer combination as internal references. Unexpectedly, relative macro and microelements content were showed a dramatic decreased compared to R₂₀:B₈₀ and the other LEDs ratios (Figure 2 and 3). While the highest levels were obtained in microgreens

were growing under R₂₀:B₈₀ combination. However, the analysis also did not take the yield into consideration.

Considering the influence of LEDs lighting combination on the microgreen's growth together with nutrition components value, we analyzed deeper the vitamin and carotenoid contents. Agreeing with our previous result obtained in the macro- and microelements, we found that the contents of Phylloquinone, α -tocopherol, Total Ascorbic Acid (TAA), and β -carotene of 21 varieties of Brassica microgreens grown under red: blue 80:20 (R₈₀:B₂₀), were significantly increased compared to other combination respectively (Figure 4).

3.2. Conclude the optimum LEDs conditions for Brassica microgreens growth conditions

Our previous data showed that LEDs lighting combination has an impact on all growth and nutritional parameters. More precisely. We found that Brassica microgreen varieties were grown under the LEDs lighting of R₇₀:G₁₀:B₂₀ combination enhances the Hypocotyl length, leaf area, fresh weight, and dry weight compared to other LEDs combination. While minerals (macro and microelements) and vitamins were significantly increased corresponding to plants grown under R₈₀:B₂₀. Attempts to detect the best LEDs combination taking into consideration the actual yield of microgreens, we conducted a correlation analysis with the yield. We estimated the minerals and vitamins concentrations corresponding to the actual fresh weight yielded (Figure 5 and 6). Interestingly, we found that mineral compositions and vitamins content in the yielded fresh weight were significantly increased in the microgreen varieties grown under the LEDs lighting of R₇₀:G₁₀:B₂₀ combination compared to other combination (Figure 5 and 6).

3.3. Hierarchical cluster analysis of 21 varieties of Brassica microgreens

A hierarchical cluster analysis profiled growth, mineral compositions and vitamins content of 21 microgreens varieties grown under R₇₀:G₁₀:B₂₀ family has been performed using *class Orange clustering hierarchical* using *ORANGE version 3.7* [29]. Presented data of microgreens grown under R₇₀:G₁₀:B₂₀, which present the highest values of growth and nutritional profile are shown in Figure 7 and Table 2. We utilized the hierarchical analysis methods (average-linkage distance between two clusters) to evaluate whether these trends were consistent across the 21 varieties under study. The hierarchical cluster analysis shows that the 21 microgreens are classified into five groups. Among the five groups, the highest distance group (Figure 7, cluster group in yellow color + Kale Tucsan in green cluster) contained 7 microgreens(Kohlrabi purple, Cabbage red, Broccoli, Kale Tucsan, Komatsuna red, Tatsoi, Cabbage green) which are representing 3 species (*B. oleracea*, *B. rapa*, *B. narinosa*).

4. Discussion

Due to the increased interest with providing the controlled environment greenhouses with LEDs lighting and for increasing the microgreen growth and nutritional profile, we investigated the impact of four different LEDs lighting ratios on the growth and nutritional quality assessment of 21 varieties belong to Brassica genera of the family Brassicaceae grew as microgreens. Microgreens are reported in many studies as valuable source vitamins, phenolics and mineral compositions [31]. Enhancing their nutritional qualities and growth is an exciting avenue of research and agriculture biotechnology.

In our study, we reveal various effects on the combination ratios between blue LEDs, red and green LEDs. A plant grown under a monochromatic light beam also stimulate specific photoreceptors that are involved in numerous biological processes. Enhance the nutritional profile and plant growth was demonstrated in many species, such as rice [32], Brassica spp. [5, 17, 22], etc. It has been reported

that red and blue light are important for the expansion of the hypocotyl elongation, pigments accumulation, and enhancement of biomass [33]. In contrast, exposure to green LEDs increases biomass at a high intensity [34]. We notice that growing microgreens under R₇₀:G₁₀:B₂₀ shows the highest value of the vegetative parameters, taking in our consideration the yield produced under all combination treatments (Figure 1). These results provide a clear indication that blue LEDs in combination with red LEDs and high-intensity green LEDs are more efficient for plants microgreen growth. Providing green lighting within the growing conditions enhance Brassica microgreens growth, while, increasing the blue light ratio had a passive response to the growth.

Many reports demonstrate the positive influence of the red:blue lighting plant growth and photosynthesis [13, 16, 17, 21, 24, 35, 36]. Furthermore, a red, blue, and green light combination has an effective source for photosynthesis [37].

Consequently, supply the red and blue LEDs combination with a green light has a significant impact on lettuce leaves growth and photosynthetic rate compared with the red and blue LEDs only. [38, 39]. It appears that blue and red light enhances the anthocyanins accumulation in leaves and become black, while green light stimulates phytochrome, shifting the active pool of Type I and Type II phytochromes to include reverse accumulation of anthocyanins [40].

Consequently, we demonstrate the positive influence of providing green light improving microgreens growth and morphology. It is reported that HL and LA of kohlrabi, mizuna, and mustard were increased when grown under green light R₇₄:G₁₈:B₈ compared with the R₈₇:B₁₃, while FW and DW were greater of those microgreens grown under providing green light than blue/red [41]. Moreover, FW of broccoli microgreens grown under light ratios of R₈₅:G₁₀:B₅ and R₈₀:B₂₀ were significantly increased than under R₇₀:G₁₀:B₂₀ [42]. The same influence is observed on chlorophyll content which improved significantly of the plant grown under additional green light [22, 41] Furthermore, the reduction on the growth parameters due to the increased of blue lighting

was reported. It is found that the hypocotyl elongation of kohrabi, mizuna, mustard was decreased under the red:blue light combination due to the inhibition of the gibberellins (GA), which inhibit the hypocotyl elongation [36].

Growing microgreens plants under blue LEDs R₂₀:B₈₀ in our study enhance the minerals composition and the vitamin content accompanied with the less growth yield compared with the green LEDs R₇₀:G₁₀:B₂₀. It is reported that broccoli microgreens grown under blue light (R₀:B₁₀₀) produce higher nutrient-dense microgreens [8, 42]. Blue light could be shown as a dominant means of regulating the nutrient content synthesis such as proton pumping, ion channel, activities, and membrane permeability [41, 42].

Comparing the LEDs lighting ratios to conclude the proper conditions, we accompanied the nutritional profile with the actual growth yielding. We found that green LEDs R₇₀:G₁₀:B₂₀ has the proper yielded influence and produced final higher mineral concentration and vitamin content due to the high growth yield. Despite the blue LED treatment to increase the mineral and vitamin content, but it is accompanied by less growth yield.

In conclusion, the assessment of 21 brassica microgreens growth and nutritional profile grown under LEDs technology provides a satisfactory growing conditions examination of microgreens. Providing green lighting ratio of R₇₀:G₁₀:B₂₀ show a positive influence on the growing microgreens growth and morphology. Interestingly, Kohlrabi purple, Cabbage red, Broccoli, Kale Tucsan, Komatsuna red, Tatsoi, Cabbage green are presented as the top microgreen's candidates of our study assessment that serve as a life support component in limited space-based conditions and controlled environment greenhouse.

Acknowledgments

Special thanks to DAAD and Exceed Swindon to provide this opportunity to present this work in the EXCEED SWINDON EXPERT WORKSHOP (Aswan-EGYPT 2018). Furthermore, Many thanks to all the lab member in the Agronomy Department, faculty of agriculture, Zagazig university for their technical support.

Author contributions. K.Y.K., A.A.N conceived and designed the experiments. K.Y.K., A.A.E, A.A.N, M.A.S.A. and S. J. L.Z. performed the experiment. K.Y.K., D.A.M., A.A.S, N.Q., S.M.A and S.Y.M analyzed the data. K.Y.K., A.A.N. wrote the manuscript. R.H., M.A.E. contributed to the manuscript writing and revision. All authors revised the manuscript.

Competing interests. The authors declare no competing financial and/or non-financial interests in relation to the work described.

Funding. This research work is a part of a project received seed funding from the Dubai Future Foundation through Gaaana.com open research platform (Grant no. MBR026).

References

- [1] B. Burlingame, Grand challenges in nutrition and environmental sustainability, *Frontiers in nutrition*, 1 (2014) 3-3.
- [2] M.C. Kyriacou, Y. Roupheal, F. Di Gioia, A. Kyrtatzis, F. Serio, M. Renna, S. De Pascale, P. Santamaria, Micro-scale vegetable production and the rise of microgreens, *Trends in Food Science & Technology*, 57 (2016) 103-115.
- [3] R. Bulgari, A. Baldi, A. Ferrante, A. Lenzi, Yield and quality of basil, Swiss chard, and rocket microgreens grown in a hydroponic system, *New Zealand Journal of Crop and Horticultural Science*, 45 (2017) 119-129.
- [4] D.D. Treadwell, R. Hochmuth, L. Landrum, W. Laughlin, Microgreen: A new specialty crop, Univ. Florida IFAS, Ext. Bul. HS1164, (2010).
- [5] Z. Xiao, G.E. Lester, Y. Luo, Q. Wang, Assessment of vitamin and carotenoid concentrations of emerging food products: edible microgreens, *J Agric Food Chem*, 60 (2012) 7644-7651.
- [6] C. Murphy, W. Pill, Cultural practices to speed the growth of microgreen arugula (*roquette*; *Eruca vesicaria* subsp. *sativa*), *The Journal of Horticultural Science and Biotechnology*, 85 (2010) 171-176.
- [7] J.S. Lee, W.G. Pill, B.B. Cobb, M. Olszewski, Seed treatments to advance greenhouse establishment of beet and chard microgreens, *The Journal of Horticultural Science and Biotechnology*, 79 (2004) 565-570.
- [8] D.A. Kopsell, C.E. Sams, Increases in Shoot Tissue Pigments, Glucosinolates, and Mineral Elements in Sprouting Broccoli after Exposure to Short-duration Blue Light from Light Emitting Diodes, *Journal of the American Society for Horticultural Science*, 138 (2013) 31-37.
- [9] M. Björkman, I. Klingen, A.N.E. Birch, A.M. Bones, T.J.A. Bruce, T.J. Johansen, R. Meadow, J. Mølmann, R. Seljåsen, L.E. Smart, D. Stewart, Phytochemicals of Brassicaceae in plant protection and human health – Influences of climate, environment and agronomic practice, *Phytochemistry*, 72 (2011) 538-556.
- [10] M. Lefsrud, D. Kopsell, A. Wenzel, J. Sheehan, Changes in kale (*Brassica oleracea* L. var. *acephala*) carotenoid and chlorophyll pigment concentrations during leaf ontogeny, *Scientia Horticulturae*, 112 (2007) 136-141.
- [11] S.D. Carvalho, K.M. Folta, Environmentally Modified Organisms – Expanding Genetic Potential with Light, *Critical Reviews in Plant Sciences*, 33 (2014) 486-508.
- [12] L. Kou, Y. Luo, T. Yang, Z. Xiao, E.R. Turner, G.E. Lester, Q. Wang, M.J. Camp, Postharvest biology, quality and shelf life of buckwheat microgreens, *LWT - Food Science and Technology*, 51 (2013) 73-78.
- [13] A. Yano, K. Fujiwara, Plant lighting system with five wavelength-band light-emitting diodes providing photon flux density and mixing ratio control, *Plant methods*, 8 (2012) 46.
- [14] N.C. Yorio, G.D. Goins, H.R. Kagie, R.M. Wheeler, J.C. Sager, Improving spinach, radish, and lettuce growth under red light-emitting diodes (LEDs) with blue light supplementation, *HortScience*, 36 (2001) 380-383.
- [15] E. Goto, Plant production in a closed plant factory with artificial lighting, *Acta Hort.*, 956 (2012) 37-49.
- [16] C. Dong, Y. Fu, G. Liu, H. Liu, Low light intensity effects on the growth, photosynthetic characteristics, antioxidant capacity, yield and quality of wheat (*Triticum aestivum* L.) at different growth stages in BLSS, *Advances in Space Research*, 53 (2014) 1557-1566.
- [17] G. Samuolienė, A. Brazaitytė, J. Jankauskienė, A. Viršilė, R. Sirtautas, A. Novičkovas, S. Sakalauskienė, J. Sakalauskaitė, P. Duchovskis, LED irradiance level affects growth and nutritional quality of Brassica microgreens, *Central European Journal of Biology*, 8 (2013) 1241-1249.
- [18] R. Matsuda, K. Ohashi-Kaneko, K. Fujiwara, E. Goto, K. Kurata, Photosynthetic characteristics of rice leaves grown under red light with or without supplemental blue light, *Plant & cell physiology*, 45 (2004) 1870-1874.

- [19] R. Matsuda, K. Ohashi-Kaneko, K. Fujiwara, K. Kurata, Effects of blue light deficiency on acclimation of light energy partitioning in PSII and CO₂ assimilation capacity to high irradiance in spinach leaves, *Plant & cell physiology*, 49 (2008) 664-670.
- [20] G.W. Stutte, S. Edney, T. Skerritt, Photoregulation of Bioprotectant Content of Red Leaf Lettuce with Light-emitting Diodes, *HortScience*, 44 (2009) 79-82.
- [21] I. Tarakanov, O. Yakovleva, I. Konovalova, G. Paliutina, A. Anisimov, Light-Emitting Diodes: on the way to combinatorial lighting technologies for basic research and crop production, *International Society for Horticultural Science (ISHS)*, Leuven, Belgium, 2012, pp. 171-178.
- [22] G. Samuolienė, A. Brazaitytė, R. Sirtautas, A. Viršilė, J. Sakalauskaitė, S. Sakalauskienė, P. Duchovskis, LED illumination affects bioactive compounds in romaine baby leaf lettuce, *Journal of the Science of Food and Agriculture*, 93 (2013) 3286-3291.
- [23] T. Ouzounis, E. Rosenqvist, C.-O. Ottosen, Spectral Effects of Artificial Light on Plant Physiology and Secondary Metabolism: A Review, *HortScience*, 50 (2015) 1128-1135.
- [24] A. Brazaitytė, S. Sakalauskienė, G. Samuolienė, J. Jankauskienė, A. Viršilė, A. Novičkovas, R. Sirtautas, J. Miliauskienė, V. Vaštakaitė, L. Dabašinskas, P. Duchovskis, The effects of LED illumination spectra and intensity on carotenoid content in Brassicaceae microgreens, *Food Chemistry*, 173 (2015) 600-606.
- [25] C.Y.L. Huang, E.E. Schulte, Digestion of plant tissue for analysis by ICP emission spectroscopy, *Communications in Soil Science and Plant Analysis*, 16 (1985) 943-958.
- [26] S.L. Booth, K.W. Davidson, J.A. Sadowski, Evaluation of an HPLC method for the determination of phyloquinone (vitamin K₁) in various food matrixes, *Journal of Agricultural and Food Chemistry*, 42 (1994) 295-300.
- [27] G.E. Lester, G.J. Hallman, J.A. Pérez, γ -Irradiation Dose: Effects on Baby-Leaf Spinach Ascorbic Acid, Carotenoids, Folate, α -Tocopherol, and Phyloquinone Concentrations, *Journal of Agricultural and Food Chemistry*, 58 (2010) 4901-4906.
- [28] D.M. Hodges, C.F. Forney, W.V. Wismer, Antioxidant Responses in Harvested Leaves of Two Cultivars of Spinach Differing in Senescence Rates, 126 (2001) 611.
- [29] J. Demsar, T. Curk, A. Erjavec, C. Gorup, T. Hocevar, M. Milutinovic, M. Mozina, M. Polajnar, M. Toplak, A. Staric, M. Stajdohar, L. Umek, L. Zagar, J. Zbontar, M. Zitnik, B. Zupan, Orange: Data Mining Toolbox in Python, *Journal of Machine Learning Research*, 14 (2013) 2349-2353.
- [30] R.G.D. Steel, J.H. Torrie, Principles and Procedures of Statistics. A Biometrical Approach, 2nd Ed. ed., MacGraw Hill Book Company, New York, 1980.
- [31] S.A. Mir, M.A. Shah, M.M. Mir, Microgreens: Production, shelf life, and bioactive components, *Crit Rev Food Sci Nutr*, 57 (2017) 2730-2736.
- [32] X.W. Chen, S.Q. Liu, Y. Wang, J.K. Liu, L. Feng, [Effects of different LED light qualities on growth, photosynthetic characteristics and nutritional quality of savoy], *Ying Yong Sheng Tai Xue Bao*, 25 (2014) 1955-1962.
- [33] H. Li, Z. Xu, C. Tang, Effect of light-emitting diodes on growth and morphogenesis of upland cotton (*Gossypium hirsutum* L.) plantlets in vitro, *Plant Cell, Tissue and Organ Culture (PCTOC)*, 103 (2010) 155-163.
- [34] M. Johkan, K. Shoji, F. Goto, S.-n. Hashida, T. Yoshihara, Blue Light-emitting Diode Light Irradiation of Seedlings Improves Seedling Quality and Growth after Transplanting in Red Leaf Lettuce, 45 (2010) 1809.
- [35] J.H. Kang, S. KrishnaKumar, S.L.S. Atulba, B.R. Jeong, S.J. Hwang, Light intensity and photoperiod influence the growth and development of hydroponically grown leaf lettuce in a closed-type plant factory system, *Horticulture, Environment, and Biotechnology*, 54 (2013) 501-509.
- [36] T.I. Potter, S.B. Rood, K.P. Zanewich, Light intensity, gibberellin content and the resolution of shoot growth in Brassica, *Planta*, 207 (1999) 505-511.

- [37] S. Muneer, E. Kim, J. Park, J. Lee, Influence of green, red and blue light emitting diodes on multiprotein complex proteins and photosynthetic activity under different light intensities in lettuce leaves (*Lactuca sativa* L.), *International journal of molecular sciences*, 15 (2014) 4657-4670.
- [38] H.H. Kim, G.D. Goins, R.M. Wheeler, J.C. Sager, Green-light supplementation for enhanced lettuce growth under red- and blue-light-emitting diodes, *HortScience*, 39 (2004) 1617-1622.
- [39] H.H. Kim, R.M. Wheeler, J.C. Sager, G.D. Gains, J.H. Naikane, EVALUATION OF LETTUCE GROWTH USING SUPPLEMENTAL GREEN LIGHT WITH RED AND BLUE LIGHT-EMITTING DIODES IN A CONTROLLED ENVIRONMENT - A REVIEW OF RESEARCH AT KENNEDY SPACE CENTER, *International Society for Horticultural Science (ISHS)*, Leuven, Belgium, 2006, pp. 111-120.
- [40] S.D. Carvalho, K.M. Folta, Green light control of anthocyanin production in microgreens, *International Society for Horticultural Science (ISHS)*, Leuven, Belgium, 2016, pp. 13-18.
- [41] J.R. Gerovac, J.K. Craver, J.K. Boldt, R.G. Lopez, Light Intensity and Quality from Sole-source Light-emitting Diodes Impact Growth, Morphology, and Nutrient Content of Brassica Microgreens, *HortScience*, 51 (2016) 497-503.
- [42] D.A. Kopsell, C.E. Sams, T.C. Barickman, R.C. Morrow, Sprouting Broccoli Accumulate Higher Concentrations of Nutritionally Important Metabolites under Narrow-band Light-emitting Diode Lighting, *Journal of the American Society for Horticultural Science*, 139 (2014) 469-477.




Figure 1. Box plot of growth and morphological measurements of *Brassica* microgreens grown under LEDs treatments. The plot illustrates the Mean and median of (Hypocotyl Length (mm), Leaf Area (cm²), Fresh weight (g), and Dry weight (%)) of 21 varieties of *Brassica* microgreens represented 5 species grown under different light-emitting diodes (LEDs) ratio (%) of red:blue 80:20 (R₈₀:B₂₀), red:blue 20:80 (R₂₀:B₈₀), red:green:blue 70:10:20 (R₇₀:G₁₀:B₂₀), or red:green:blue 20:10:20 (R₂₀:G₁₀:B₇₀) (Supplemental Table 2 and 3). Resulting ranking could be analyzed with point values of Mean and Median or uncertainty range with box. Statistical analysis is performed using a one-way ANOVA test (P ≤ 0.05). Small letters denote statistically significant differences.




Figure 2. Box plot of mineral composition and content of macroelements of *Brassica* microgreens grown under LEDs treatments. The plot illustrates the Mean and median of macroelements concentrations; P, K, Ca and Na (mg/100 g FW) of 21 varieties of *Brassica* microgreens represented 5 species grown under different light-emitting diodes (LEDs) ratio (%) of red:blue 80:20 (R₈₀:B₂₀), red:blue 20:80 (R₂₀:B₈₀), red:green:blue 70:10:20 (R₇₀:G₁₀:B₂₀), or red:green:blue 20:10:20 (R₂₀:G₁₀:B₇₀) (Supplemental Table 4 and 5). Resulting ranking could be analyzed with point values of Mean and Median or uncertainty range with box. Statistical analysis is performed using a one-way ANOVA test (P ≤ 0.05). Small letters denote statistically significant differences.




Figure 3. Box plot of mineral composition and content of microelements of *Brassica* microgreens grown under LEDs treatments. The plot illustrates the Mean and median of microelements concentrations; Fe, Zn, Cu and Mn (mg/100 g FW) of 21 varieties of *Brassica* microgreens represented 5 species grown under different light-emitting diodes (LEDs) ratio (%) of red:blue 80:20 (R₈₀:B₂₀), red:blue 20:80 (R₂₀:B₈₀), red:green:blue 70:10:20 (R₇₀:G₁₀:B₂₀), or red:green:blue 20:10:20 (R₂₀:G₁₀:B₇₀) (Supplemental Table 6 and 7). Resulting ranking could be analyzed with point values of Mean and Median or uncertainty range with box. Statistical analysis is performed using a one-way ANOVA test ($P \leq 0.05$). Small letters denote statistically significant differences.




Figure 4. Box plot of vitamin and carotenoid concentrations of *Brassica* microgreens grown under LEDs treatments. The plot illustrates the Mean and median of vitamin and carotenoids concentrations; Phylloquinone (ug/ g FW), α-tocopherol, Total Ascorbic Acid (TAA), and β-carotene (mg/100 g FW) of 21 varieties of Brassica microgreens represented 5 species grown under different light-emitting diodes (LEDs) ratio (%) of red:blue 80:20 (R₈₀:B₂₀), red:blue 20:80 (R₂₀:B₈₀), red:green:blue 70:10:20 (R₇₀:G₁₀:B₂₀), or red:green:blue 20:10:20 (R₂₀:G₁₀:B₇₀) (Supplemental Table 8 and 9) . Resulting ranking could be analyzed with point values of Mean and Median or uncertainty range with box. Statistical analysis is performed using a one-way ANOVA test ($P \leq 0.05$). Small letters denote statistically significant differences.




Figure 5. Mineral composition and content of *Brassica* microgreens under LEDs treatments. A) Mean macroelement concentration of P, K, Ca, and Na. **B)** Mean microelement concentration of Fe, Zn, Cu, and Mn of 21 varieties of *Brassica* microgreens exposed to different light-emitting diodes (LEDs) ratio (%) of red:blue 80:20 (R₈₀:B₂₀), red:blue 20:80 (R₂₀:B₈₀), red:green:blue 70:10:20 (R₇₀:G₁₀:B₂₀), or red:green:blue 20:10:20 (R₂₀:G₁₀:B₇₀). Data represents as a mean concentration corresponding to the actual Fresh weight (fresh weight results of each LEDs treatments of the 21 varieties (Supplementary Table 3) (Actual concentration (mg/ g actual FW) = Concentration (mg/ 100 g FW) X Fresh weight (g) / 100). Mean±SE values are based on a representative sample from each treatment across three experimental replications. * for significant at $P \leq 0.05$.




Figure 6. Assessment of vitamin and carotenoid concentrations of *Brassica* microgreens under LEDs treatments. A) Mean α -tocopherol, Total Ascorbic Acid (TAA), and β -carotene (mg/100 g FW) concentration. **B)** Mean Phylloquinone (μ g/ g FW) concentration of 21 varieties of *Brassica* microgreens exposed to different light-emitting diodes (LEDs) ratio (%) of red:blue 80:20 (R₈₀:B₂₀), red:blue 20:80 (R₂₀:B₈₀), red:green:blue 70:10:20 (R₇₀:G₁₀:B₂₀), or red:green:blue 20:10:20 (R₂₀:G₁₀:B₇₀). Data represents as a mean concentration corresponding to the actual Fresh weight (fresh weight results of each LEDs treatments of the 21 varieties (supplementary Table 3) (Actual concentration (mg/ g actual FW) = Concentration (mg/ 100 g FW) X Fresh weight (g) / 100). For Phylloquinone ((Actual concentration (mg / g actual FW) = Concentration (μ g/ g FW) X Fresh weight (g)). Mean±SE values are based on a representative sample from each treatment across three experimental replications. * for significant at $P \leq 0.05$.




Figure 7. The average-linkage on the normalized data sets of mineral composition and vitamin and carotenoid concentrations corresponding to the actual fresh weight by means of the Hierarchical method using growth and morphology measurements data of 21 varieties Brassica microgreens grown under light-emitting diodes (LEDs) ratio (%) of red:green:blue 70:10:20 ($R_{70}:G_{10}:B_{20}$). The complete profile of the highest cluster value (Yellow cluster) microgreens presented in Table 2.

Tables

Table 1. Twenty-one varieties of Brassica microgreens represented 5 species Brassica genera assayed in this study. Growth length (day) and seed index (g) show each variety growth period and the number of seeds used for the sowing.

Commercial name	Scientific name (genus and species)	Growth length (day)	Seed index (g)
Broccoli	<i>Brassica oleracea</i> L. var. <i>italica</i>	9	10
Brussel sprouts	<i>Brassica oleracea</i> L. var. <i>Gemmifera</i>	10	15
Cabbage green	<i>Brassica oleracea</i> L. var. <i>capitata</i> f. <i>alba</i>	9	10
Cabbage red	<i>Brassica oleracea</i> L. var. <i>capitata</i> f. <i>rubra</i>	8	10
Cabbage savoy	<i>Brassica oleracea</i> L. var. <i>capitata</i> f. <i>sabauda</i>	8	10
Cauliflower	<i>Brassica oleracea</i> L. var. <i>botrytis</i>	9	15
Collard	<i>Brassica oleracea</i> L. var. <i>viridis</i>	10	15
Kale Chinese	<i>Brassica oleracea</i> L. var. <i>alboglabra</i>	10	15
Kale red	<i>Brassica oleracea</i> L. var. <i>acephala</i>	9	10
Kale Tucsan	<i>Brassica oleracea</i> L. var. <i>acephala</i>	9	15
Kohlrabi purple	<i>Brassica oleracea</i> L. var. <i>gongylodes</i>	7	25
Cabbage Chinese	<i>Brassica rapa</i> L. var. <i>pekinensis</i>	6	15
Komatsuna red	<i>Brassica rapa</i> L. var. <i>perviridis</i>	8	15
Mizuna	<i>Brassica rapa</i> L. var. <i>nipposinica</i>	8	15
Pak choy	<i>Brassica rapa</i> L. var. <i>chinensis</i>	8	15
Rapini	<i>Brassica rapa</i> L. var. <i>ruvo</i>	9	15
Turnip	<i>Brassica rapa</i> L. var. <i>rapa</i>	9	10
Mustard Dijon	<i>Brassica juncea</i> (L.) Czern.	12	15
Mustard red	<i>Brassica juncea</i> (L.) Czern.	10	10
Rutabaga	<i>Brassica napus</i> L. var. <i>napobrassica</i>	9	10
Tatsoi	<i>Brassica narinosa</i> L. var. <i>rosularis</i>	7	10

Table 2. Growth, and nutritional composition profile of highest Brassica microgreens grown under light-emitting diodes (LEDs) ratio (%) of red:green:blue 70:10:20 (R₇₀:G₁₀:B₂₀). List of the 7 brassica microgreens is exported from the hierarchical cluster analysis (Figure 7).

	Kohlrabi purple	Cabbage red	Broccoli	Kale Tucson	Komatsuna red	Tatsoi	Cabbage green
Hypocotyl Length (mm)	46	47	42	49	44	46	45
Leaf Area (cm ²)	1.82	1.93	1.65	2.02	1.75	1.86	1.83
Fresh weight (g)	0.44	0.46	0.42	0.47	0.41	0.45	0.44
Dry weight (%)	6.65	6.18	6.55	6.90	6.57	6.34	6.25
P (mg/100 g FW)	68	62	59	63	66	64	62
K (mg/100 g FW)	322	224	319	280	320	300	183
Ca (mg/100 g FW)	65	84	92	55	53	44	87
Na (mg/100 g FW)	46	40	50	46	28	35	69
Fe (mg/100 g FW)	0.77	0.69	0.74	0.76	0.76	0.65	0.67
Zn (mg/100 g FW)	0.43	0.40	0.42	0.38	0.38	0.41	0.33
Cu (mg/100 g FW)	0.08	0.10	0.11	0.07	0.05	0.08	0.06
Mn (mg/100 g FW)	0.39	0.35	0.41	0.46	0.34	0.35	0.37
Phylloquinone (ug/ g FW)	2.6	2.3	2.0	1.7	2.2	1.5	1.4
α-tocopherol (mg/100 g FW)	17.6	29.5	17.3	19.4	25.0	29.3	14.3
Total Ascorbic Acid (mg/100 g FW)	77.1	127.4	84.6	73.2	97.5	99.5	118.9
β-carotene (mg/100 g FW)	6.6	9.9	6.9	5.4	7.1	10.6	9.6